

RESTRICTED USE PESTICIDE
DUE TO POTENTIAL SECONDARY TOXICITY TO NONTARGET ORGANISMS

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF IDAHO

rozol[®] VOLE BAIT

FOR AERIAL BROADCAST APPLICATION

FOR CONTROL OF VOLES (*Microtus* spp.) OVER POME AND STONE FRUIT ORCHARDS, NURSERIES, TREE AND FORESTRY PLANTATIONS, CHRISTMAS TREE FARMS, AND FALLOW AGRICULTURAL LAND, OR WITHIN 100 FEET OF THE EDGE OF CROPLAND ON RANGELAND AND NONCROP AREAS INCLUDING BORDER AREAS, BUFFER STRIPS AND CENTER PIVOT CORNERS ADJACENT TO CROPS

EPA SLN No. ID-190004

EPA Reg. No. 7173-242

This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2024.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions and precautions on the EPA registered label.

DIRECTIONS FOR USE:

This product may be applied by broadcast from aircraft, for control of voles (*Microtus* spp.) over pome and stone fruit orchards, nurseries, tree and forestry plantations, Christmas tree farms, and fallow agricultural land, or within 100 feet of the edge of cropland on rangeland and noncrop areas including border areas, buffer strips and center pivot corners adjacent to crops. This product must not be applied directly to food or feed crops except as specified on this label.

APPLICATION DIRECTIONS:

Before application, locate vole trails, runway systems and harborage areas to be treated. Uniformly apply 10 pounds of bait per acre using commercial bait dispensing equipment. If populations are high, make a second application 1 to 2 months after the first.

RESTRICTIONS/PRECAUTIONS:

- Do not apply more than 10 pounds of bait per acre per application.
- Do not apply more than 20 pounds of bait per acre per season.
- In pome and stone fruit orchards, apply bait after all fruit is harvested (including drops) and before new growth in the spring. Do not apply bait during the growing season.
- Apply bait when no rain or snow is expected for three consecutive days.
- Do not allow animals to graze in treated areas.
- Keep away from humans, domestic animals and pets during application, handling and storage.
- Do not apply bait in a manner that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Keep all other persons out of the treated area during application.
- Do not apply within 50 feet of any body of surface water or where raptors are actively feeding on voles.
- This product is extremely toxic to mammals and birds. This SLN label should not be used where impact on listed threatened or endangered species is likely. Consult the Federal label for additional restrictions and precautions to protect fish and wildlife.

24(c) registrant:

Liphatech, Inc.
3600 W. Elm Street
Milwaukee, WI 53209
(800) 351-1476

LIPHATECH[®]

04/16/2019