

# RESTRICTED USE PESTICIDE

## DUE TO HAZARD TO NONTARGET ORGANISMS

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certificate.

## Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

# rozol<sup>®</sup> VOLE BAIT

FOR AERIAL BROADCAST APPLICATION TO CONTROL MEADOW OR MOUNTAIN VOLES  
IN POME FRUIT (APPLE, PEAR) AND STONE FRUIT (APRICOT, CHERRY, NECTARINE,  
PEACH, PLUM, PRUNE) TREE ORCHARDS AFTER FALL HARVEST

EPA Reg. No. 7173-242 EPA SLN No. WA-070019

This label for Rozol Vole Bait expires and must not be distributed or used in accordance with this SLN registration after December 31, 2024.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions and precautions on this SLN label and the EPA registered label.

### DIRECTIONS FOR USE:

#### POME AND STONE FRUIT TREE ORCHARDS FOR MEADOW OR MOUNTAIN VOLE CONTROL:

**APPLICATION DIRECTIONS:** Before application, examine orchard floor to locate trails and runway systems to be treated. Apply bait when three days of rain-free and snow-free weather are expected. Apply bait after harvest (including all harvestable fruit drops) and before new growth in the spring.

**Aerial broadcast baiting:** Uniformly apply 10 pounds of bait per acre. If populations are high, make a second application 1 to 2 months after the first. Do not apply more than 20 pounds of bait per acre per season. To avoid exposing nontarget organisms, clean up any bait in piles for reuse or disposal according to the federal label.

#### RESTRICTIONS / PRECAUTIONS:

- Do not apply more than 10 pounds of bait per acre per application.
- Do not apply more than 20 pounds of bait per acre per season.
- Do not apply bait during the growing season.
- Do not apply bait over bare ground or to crops not specified above.
- Do not expose humans (including children), pets, livestock or other nontarget animals to bait during application, handling, storage or disposal. Do not allow livestock to graze or clean up fruit drops in treated areas. Do not use hay cut after application for livestock feed or bedding.
- Do not apply this product directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.
- Do not apply by air within 50 feet of any body of surface water (i.e. streams, rivers, lakes) or where raptors are actively feeding on voles.
- This pesticide is extremely toxic to fish and wildlife. Rozol Vole Bait should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may refer to the Natural Resources Assessment Section on WSDA's website for additional information on pesticide use and endangered species protection. You may use the Washington Department of Fish and Wildlife's online SalmonScape mapping tool to determine if listed salmonid species occur in or near a proposed application site.

#### WSDA Container Disposal Guidance:

Completely empty the contents of the container into application equipment, and apply to a labeled crop or site. The empty container may be disposed of in a sanitary landfill. Burning is not a legal method of container disposal in Washington.

Rev. 11/01/2023

Section 24(c) SLN registrant:  
Liphatech, Inc.  
3600 W. Elm Street  
Milwaukee, WI 53209

**LIPHATECH<sup>®</sup>**

11/01/23